

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:
75-684

APPROVED DRAFT LABELING

Note: Keyline does not print.

FAMOTIDINE INJECTION

**PHARMACY BULK PACKAGE -
NOT FOR DIRECT INFUSION**
MUST BE DILUTED PRIOR TO IV USE.
Do not dispense as a unit.

500 mg/50 mL *

20 mg/2 mL
Rx ONLY.

NDC 55390-026-01

50 mL PHARMACY BULK VIAL
FOR THE PREPARATION OF INTRAVENOUS SOLUTIONS AND USUAL ADULT DOSAGE:
See package insert.

*Each mL contains 10 mg of famotidine and the following inactive ingredients: L-aspartic acid 4 mg, mannitol 20 mg, and Water for Injection, q.s., 1 mL. Benzyl alcohol 9 mg added as preservative.

Store at 2° to 8°C (36° to 46°F).

Once the container closure has been punctured, withdrawal of the container contents should be completed without delay. THE ENTIRE CONTENTS OF THE VIAL SHOULD BE DISPENSED WITHIN 4 HOURS OF INITIAL ENTRY.

Date Entered: _____

Time of Entry: _____

Manufactured by:
Ben Venue Labs, Inc.
Bedford, OH 44146

BEDFORD
LABORATORIES

Manufactured for:
Bedford Laboratories™
Bedford, OH 44146

FAMP-P8V008

LOT
EXP

APR 16 2001

PRINTED SIDE
C011294HA

1-1-1

+

*Each mL contains 10 mg of famotidine and the following inactive ingredients: L-aspartic acid 4 mg, mannitol 20 mg, and Water for Injection, q.s., 1 mL. Benzyl-alcohol 9 mg added as preservative.

APR 16 2001
APPROVED

Manufactured by:
Ben Venue Labs, Inc.
Bedford, OH 44146

Manufactured for:
Bedford Laboratories™
Bedford, OH 44146

NDC 55390-026-01
50 mL PHARMACY BULK VIAL

FAMOTIDINE INJECTION

PHARMACY BULK PACKAGE—
NOT FOR DIRECT INFUSION

MUST BE DILUTED PRIOR TO IV USE.
Do not dispense as a unit.

500 mg/50 mL*

20 mg/2 mL
Rx ONLY.

BEDEORD
LABORATORIES™



LOT

EXP

FAM-PBC00C

FOR THE PREPARATION OF
INTRAVENOUS SOLUTIONS
and USUAL ADULT DOSAGE:
See accompanying circular.

Store at 2° to 8°C (36° to 46°F).

If solution freezes, bring to
room temperature; allow
sufficient time to solubilize.

Once the container closure
has been punctured, withdrawal
of the container contents should
be completed without delay. THE
ENTIRE CONTENTS OF THE
VIAL SHOULD BE
DISPENSED WITHIN 4 HOURS
OF INITIAL ENTRY.

20 mg/2 mL
500 mg/50 mL*
PHARMACY BULK PACKAGE—
NOT FOR DIRECT INFUSION

FAMOTIDINE INJECTION

NDC 55390-026-01
50 mL PHARMACY BULK VIAL

FAMOTIDINE INJECTION

PHARMACY BULK PACKAGE—
NOT FOR DIRECT INFUSION

MUST BE DILUTED PRIOR TO IV USE.
Do not dispense as a unit.

500 mg/50 mL*

20 mg/2 mL
Rx ONLY.

BEDEORD
LABORATORIES™

1-1-1

+

Format Number: 77369 #086A
Black
478 Brown
032 Red

Prepared by
Mark Zarnstorff
Checked by

